

Used as directed at 200 to 400 ppm concentration in waters up to 500 ppm of hardness calculated as CaCO₃, this product is effective against the following pathogenic organisms in 60 seconds:

Community Associated Methicillin Resistant *Staphylococcus Aureus* (CA-MRSA), *Escherichia coli*, *Escherichia coli* 0157:H7, HIV-1 (AIDS Virus), *Listeria monocytogenes*, *Staphylococcus aureus*, *Shigella dysenteriae*, and *Yersinia enterocolitica*.

STERAMINE sanitizing solutions of 200 to 400 ppm concentration may be used on food processing equipment, utensils and other food contact articles and surfaces as specified under 40 CFR 180.940 of the US Environmental Protection Agency.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS and DOMESTIC ANIMALS**

WARNING: Tablet dust causes substantial but temporary eye injury. Do not get dust in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Tablets may be harmful if swallowed.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN: Rinse immediately with water for 15-20 minutes. If irritation persists, call poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by poison control center doctor. Do not give anything by mouth to an unconscious person. Have the product label or container with you when calling a poison control center or doctor, or going for treatment.

STORAGE AND DISPOSAL. Do not contaminate water, food or feed by storage and disposal. Store in original container in areas inaccessible to persons unfamiliar with its use. Nonrefillable container. Do not reuse or refill this container. Triple rinse promptly after emptying. Offer empty container for recycling if available or discard in trash. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

S11-1011

www.sanitize.com

STERAMINE™

1-G Tablets

The Multi-Purpose Sanitizer



FOR SANITIZING FOOD CONTACT SURFACES

USE ONE TO TWO TABLETS PER 1 GALLON OF WATER

Kills HIV-1 (AIDS Virus) and Community Associated Methicillin Resistant *Staphylococcus aureus* (CA-MRSA) when used as directed for sanitizing.

EPA REG. No. 1561-11 EPA EST. No. 1561-VA-1

ACTIVE INGREDIENT: (Quaternary) Alkyl (C14 95%. C12 3%. C16 2%) dimethyl benzyl ammonium chloride dihydrate	50%
OTHER INGREDIENTS	50%
TOTAL	100%

KEEP OUT OF REACH OF CHILDREN WARNING

See left panel for additional precautionary statements.
For Institutional and Commercial use only.

NET WEIGHT
8 OUNCES

150
Tablets

Manufactured by
EDWARDS-COUNCILOR CO. INC., 1427 Baker Road, Virginia Beach, VA 23455

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Prepare sanitizing solutions with warm water. Allow several minutes for tablets to dissolve before using.

FOR SANITIZING DISHES, GLASSES, AND UTENSILS IN RESTAURANTS, TAVERNS, AND OTHER PUBLIC EATING PLACES.

1. Scrape and prewash utensils and glasses whenever possible.
2. Wash with a good detergent or compatible cleaner in first sink compartment.
3. Rinse with clean water in second sink compartment.
4. Sanitize in a solution of 1 to 2 TABLETS per 1 GALLON OF WATER (200 to 400 ppm) in third sink compartment. Immerse all utensils for at least one minute or for contact time specified by governing sanitary code.
5. Place sanitized utensils on a rack or drainboard to air dry.
6. A fresh sanitizing solution must be prepared at least daily or more often if the solution becomes diluted or soiled.

DIRECTIONS FOR SPRAYING

FOR SANITIZING FOOD PROCESSING EQUIPMENT, DAIRY EQUIPMENT, SINKS, COUNTERTOPS, REFRIGERATED STORAGE AND DISPLAY EQUIPMENT AND OTHER HARD NON-POROUS FOOD CONTACT ARTICLES AND SURFACES.

1. Wash and rinse all articles and surfaces thoroughly.
2. Apply a solution of 200 to 400 ppm concentration by combining 1 or 2 tablets per gallon of warm water, allowing several minutes for tablets to dissolve completely before using. Spray with hand trigger sprayer or wipe on and allow surface to remain wet for at least one minute followed by adequate draining and air drying.

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS

Personal Protection: The specific barrier protection items to be used when handling items soiled with blood or body fluids are disposable gloves, gowns, masks and eye coverings.

Cleaning Procedure: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

Contact Time: One minute.

Disposal of Infectious Materials: Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal.

